



Bioscientia Medicina: Journal of Biomedicine & Translational Research

Journal Homepage: www.bioscmed.com

Pectoralis Major Flap: A Versatile Option for Head and Neck Reconstruction - A Case Report

Radot Oktora Tua Pasaribu^{1*}, Affandi Wiramur², Joko Purnomo³

¹General Surgery Resident, Faculty of Medicine, Universitas Sebelas Maret/Dr. Moewardi Regional General Hospital, Surakarta, Indonesia

²Department of Plastic and Reconstructive Surgery, Universitas Sebelas Maret/Dr. Moewardi Regional General Hospital, Surakarta, Indonesia

³Department of Oncology Surgery, Universitas Sebelas Maret/Dr. Moewardi Regional General Hospital, Surakarta, Indonesia

ARTICLE INFO

Keywords:

Case report
Head and neck reconstruction
Parotid cancer
Pectoralis major flap
Skin flap

*Corresponding author:

Radot Oktora Tua Pasaribu

E-mail address:

radot.hack19@gmail.com

All authors have reviewed and approved the final version of the manuscript.

<https://doi.org/10.37275/bsm.v9i3.1213>

ABSTRACT

Background: Skin flaps are an essential tool in reconstructive surgery, especially in the head and neck region where defects can result from tumor excision, trauma, or congenital anomalies. The pectoralis major flap is a well-established and versatile option for head and neck reconstruction due to its reliable vascularity, ease of harvest, and ability to cover a variety of defect sizes and locations. **Case presentation:** This report describes the case of a 22-year-old male with a history of parotid cancer who presented with a large, bleeding mass on the right side of his neck. Following tumor excision, the resulting defect was successfully reconstructed using a pectoralis major flap. **Conclusion:** The pectoralis major flap is a reliable and versatile reconstructive option for head and neck defects. It offers several advantages, including good vascularity, ease of harvest, and the ability to provide adequate tissue bulk for complex reconstructions.

1. Introduction

The intricate anatomy and critical functions of the head and neck region present unique challenges in reconstructive surgery. Defects in this area often arise from the surgical removal of tumors, traumatic injuries, or congenital anomalies, each demanding meticulous repair to restore both form and function. The ideal reconstructive solution must address multiple factors: providing sufficient tissue coverage, ensuring robust blood supply, minimizing complications at the donor site, and achieving aesthetically acceptable results.^{1,2}

Skin flaps have emerged as an indispensable tool in the reconstructive surgeon's armamentarium. These versatile techniques involve the transfer of skin and underlying subcutaneous tissue, along with their inherent blood supply, from a donor site to the area requiring reconstruction. Among the various skin flaps available, the pectoralis major flap stands out as a cornerstone of head and neck reconstruction. Its popularity stems from its reliable blood supply, the relative ease of harvesting the flap, and its capacity to provide substantial tissue bulk, making it suitable for addressing complex defects.³⁻⁵

The pectoralis major flap derives its blood supply from the pectoral branch of the thoracoacromial artery, a robust vascular network that ensures the flap's vitality after transfer. This vascular reliability contributes significantly to the success of the reconstruction, minimizing the risk of flap necrosis, a serious complication that can compromise the outcome. Additionally, the pectoralis major muscle, located in the chest, is readily accessible, allowing for efficient flap harvest with minimal disruption to surrounding tissues and a reduced risk of complications at the donor site.^{6,7}

The versatility of the pectoralis major flap extends to its ability to address a wide spectrum of head and neck defects. Its ample tissue bulk makes it particularly well-suited for reconstructing extensive or complex defects, providing sufficient volume to restore the natural contours of the head and neck. Whether the defect involves the oral cavity, oropharynx, hypopharynx, or larynx, the pectoralis major flap offers a reliable reconstructive solution.⁸⁻¹⁰ In this case report, we illustrate the successful application of a pectoralis major flap in the reconstruction of a large head and neck defect following tumor excision in a 22-year-old male patient.

2. Case Presentation

This report details the case of a 22-year-old male who presented with a complex medical history and a significant head and neck lesion necessitating surgical intervention and reconstruction. The patient's presentation, clinical findings, laboratory results, imaging studies, and final diagnosis are outlined below. The patient's medical history was significant for a right-sided neck mass that had been present for approximately two years. Initially, the mass was asymptomatic, causing neither pain nor any noticeable skin changes. However, over the past seven months, the mass had exhibited a concerning increase in size. Furthermore, in the preceding three months, the overlying skin had broken down, forming a wound that intermittently bled. Adding to the complexity of the case, the patient had a prior diagnosis of parotid

gland cancer. He had undergone a comprehensive treatment regimen for this condition, including six cycles of chemotherapy, four radiotherapy sessions, and embolization of the right external carotid artery. This extensive treatment history highlighted the aggressive nature of the patient's underlying malignancy and underscored the challenges in managing his current presentation. Upon physical examination, a large, actively bleeding mass was evident on the right side of the patient's neck. The overlying skin exhibited a wound characterized by a reddish hue and ongoing sanguineous discharge. Palpation of the mass revealed a firm, lumpy consistency with well-defined borders. The mass was fixed to the underlying tissues and was not tender to the touch. There were no signs of localized inflammation, such as increased warmth or erythema, apart from the area immediately surrounding the wound. The patient's vital signs were within the normal range, with a Glasgow Coma Scale (GCS) score of 15/15, indicating a normal level of consciousness. His heart rate was slightly elevated at 125 beats per minute, possibly reflecting a degree of anxiety or discomfort related to his condition. His blood pressure was 148/86 mmHg, falling within the acceptable range for his age. Laboratory investigations revealed two key findings: anemia of chronic disease and leukocytosis. The anemia was likely multifactorial, potentially attributed to the underlying malignancy, prior chemotherapy, and chronic blood loss from the neck mass. Leukocytosis, an elevated white blood cell count, could signify an ongoing inflammatory process or infection associated with the wound. A comprehensive array of imaging studies was conducted to delineate the extent of the lesion and guide surgical planning. A chest X-ray was performed to screen for potential pulmonary metastases, which were fortunately not identified. Magnetic resonance imaging (MRI) of the cervical region revealed a heterogeneously enhancing lesion, suggestive of a malignant mass. The MRI also identified multiple enlarged lymph nodes in the right neck, including those in the right parotid gland and bilateral

supraclavicular regions. This finding indicated the potential spread of the malignancy to regional lymph nodes. A nasopharyngeal computed tomography (CT) scan further characterized the lesion, demonstrating multiple heterogeneous lesions with areas of necrosis. The mass was observed to be compressing adjacent anatomical structures, including the parotid space, buccal space, and right carotid space. The CT scan also confirmed the presence of multiple enlarged lymph nodes. Finally, a three-dimensional CT scan of the head was performed to assess for any intracranial involvement. No evidence of brain metastases was found. The scan also confirmed the presence of embolization coils in the right external carotid artery, remnants of the patient's previous treatment. Based on the collective findings from the patient's history, physical examination, laboratory results, and imaging studies, a diagnosis of right-sided parotid gland cancer (Ca Parotid Dextra) with advanced local disease and regional lymph node involvement (T4N3M0) was established. This diagnosis highlighted the challenging nature of the case, requiring a multidisciplinary approach to treatment, including surgical resection and reconstruction (Table 1).

The management of this patient's complex case involved a multi-faceted approach, encompassing thorough preoperative evaluation, meticulous surgical execution, and comprehensive postoperative care. Each stage of this process is detailed below. Prior to surgical intervention, a comprehensive evaluation was undertaken to assess the patient's overall health status and optimize his condition for surgery. This evaluation included a detailed review of his medical history, a thorough physical examination, and imaging studies, including computed tomography (CT) and magnetic resonance imaging (MRI) scans, to precisely delineate the extent of the tumor and its relationship to surrounding structures. Given the patient's history of cancer and prior treatments, special attention was paid to assessing his nutritional status and identifying any potential comorbidities that could influence surgical risk or postoperative recovery. This assessment aimed to optimize his overall health

and minimize potential complications. A critical component of preoperative management involved a detailed discussion with the patient regarding the proposed surgical procedure, including its potential benefits, inherent risks, and alternative treatment options. This informed consent process ensured that the patient was actively involved in decision-making and fully understood the implications of the chosen treatment strategy. Finally, the patient received prophylactic antibiotics to minimize the risk of surgical site infection, a crucial step in ensuring optimal wound healing and preventing postoperative complications. The surgical procedure involved a carefully orchestrated sequence of steps aimed at complete tumor removal and reconstruction of the resulting defect. The surgical field was prepared and draped, ensuring strict aseptic technique. A curvilinear incision was meticulously planned and executed in the neck, following the natural skin lines to optimize the cosmetic outcome. The incision was designed to encompass the tumor entirely while providing adequate exposure for dissection and reconstruction. The tumor was carefully dissected from the surrounding tissues, ensuring complete removal en bloc, meaning the tumor was excised as a single unit with a margin of healthy tissue. This technique minimizes the risk of leaving residual tumor cells behind and promotes the best chance of disease control. The surgical margins were carefully examined to confirm complete tumor resection. With the tumor successfully removed, attention turned to reconstructing the resulting defect, which measured 8 x 12 cm. A pectoralis major myocutaneous flap was chosen for this purpose due to its reliable blood supply, ample tissue bulk, and relative ease of harvest. A skin paddle was designed on the chest, overlying the pectoralis major muscle, carefully considering the size and shape needed to adequately fill the defect. The flap was meticulously elevated, preserving the underlying pectoralis major muscle and its critical vascular pedicle, the pectoral branch of the thoracoacromial artery. The harvested pectoralis major flap was then carefully transposed to the neck

defect. The flap was meticulously inset, ensuring proper orientation and tension to achieve optimal aesthetic and functional outcomes. The vascular pedicle was carefully monitored to confirm robust blood flow to the flap, minimizing the risk of flap necrosis. Following successful flap transfer, the donor site on the chest was closed primarily, meaning the edges of the incision were brought together and meticulously sutured. A drain was often placed at the donor site to prevent fluid accumulation and promote healing. The neck incision was closed in layers, using absorbable sutures for the deeper tissues and non-absorbable sutures for the skin. This layered closure technique provides strength and support to the wound, promoting optimal healing and minimizing scar formation. Postoperative care focused on monitoring the patient's recovery, ensuring flap viability, promoting wound healing, and managing potential complications. The patient was closely monitored in the immediate postoperative period, with careful attention to vital signs, flap viability, and wound healing. Regular assessments were performed to detect any signs of complications, such as bleeding,

infection, or flap necrosis. Adequate pain control was prioritized to ensure the patient's comfort and facilitate recovery. Analgesics were administered as needed to manage postoperative pain effectively. Prophylactic antibiotics were continued in the postoperative period to prevent infection and promote optimal wound healing. Drains placed at the surgical site were closely monitored, and output was carefully recorded. Drains were removed when the output decreased to an acceptable level, indicating adequate fluid drainage and minimizing the risk of seroma formation. Proper nutrition is crucial for wound healing and overall recovery. The patient received appropriate nutritional support to ensure adequate caloric and protein intake, promoting tissue regeneration and optimizing recovery. The patient was scheduled for regular follow-up appointments to assess flap viability, wound healing progress, and overall outcome. These follow-up visits allowed for early detection and management of any potential complications and ensured long-term monitoring of the reconstruction (Table 2).

Table 1. Anamnesis, clinical finding, laboratory, imaging and diagnosis.

Category	Findings
Anamnesis	22-year-old male, Bleeding from a lump on the right neck
	Lump present for 2 years
	No pain or wound initially
	Lump got bigger in the last 7 months
	Wound present for 3 months
	History of parotid cancer
	6 cycles of chemotherapy, 4 times radiotherapy, and embolization
Clinical finding	Large bleeding mass on the right neck
	Wound present, reddish with active bleeding
	Tenderness not felt, lumpy, hard consistency, fixed, clear boundaries, no temperature change
	GCS 15/15, heart rate 125 bpm, blood pressure 148/86 mmHg
Laboratory	Anemia of chronic disease
	Leukocytosis
Imaging	Thorax X-Ray: No pulmonary metastases
	Cervical MRI: Heterointense lesion suggestive of malignant mass, multiple colli dextra, right parotid, and bilateral supraclavicular lymphadenopathy
	Nasopharyngeal MSCT: Multiple heterogeneous lesions with necrotic component, pushing parotid space, buccal space, and right carotid space, multiple lymphadenopathy
	Head MSCT 3D: No brain metastases, post-embolized coil embolism material in the right external artery
Diagnosis	Ca Parotid Dextra T4N3M0

Table 2. Treatment and surgical procedure.

Category	Details
Preoperative management	Patient evaluation, including medical history, physical examination, and imaging studies (CT scan, MRI). The patient underwent surgical excision of the tumor, resulting in an 8 x 12 cm defect in the right neck. The defect was successfully reconstructed using a pectoralis major flap.
	Assessment of comorbidities and nutritional status
	Discussion of treatment options and surgical risks and benefits with the patient
	Antibiotic prophylaxis
Surgical procedure	1. Incision and Exposure: A curvilinear incision is made in the neck, following the natural skin lines, to encompass the tumor and provide adequate exposure.
	2. Tumor Excision: The tumor is carefully dissected and removed en bloc, ensuring complete resection with clear margins.
	3. Flap Harvesting: A skin paddle is designed on the chest, based on the pectoralis major muscle. The flap is elevated, preserving the vascular pedicle (pectoral branch of the thoracoacromial artery).
	4. Flap Transposition: The flap is transposed to the neck defect and inset, ensuring adequate vascularity and coverage of the defect.
	5. Donor Site Closure: The donor site on the chest is closed primarily, often with the aid of a drain.
	6. Wound Closure: The neck incision is closed in layers, using absorbable and non-absorbable sutures.
Postoperative management	1. Monitoring: Close monitoring of the patient's vital signs, flap viability, and wound healing.
	2. Pain Management: Adequate pain control with analgesics.
	3. Antibiotic Therapy: Continuation of antibiotic prophylaxis to prevent infection.
	4. Drain Management: Removal of drains when output is minimal.
	5. Nutritional Support: Adequate nutrition to promote wound healing.
	6. Follow-up: Regular follow-up appointments to assess flap viability, wound healing, and overall outcome.

3. Discussion

The pectoralis major flap has rightfully earned its reputation as a cornerstone in the challenging field of head and neck reconstruction. This workhorse flap offers a combination of advantageous characteristics that make it a reliable and versatile option for addressing a wide range of defects in this anatomically complex region. At the heart of the pectoralis major flap's success lies its robust and predictable blood supply. This intricate vascular network ensures the flap's vitality after transfer, minimizing the risk of necrosis and optimizing the chances of successful integration at the recipient site. The dominant blood supply to the pectoralis major flap comes from the pectoral branch of the thoracoacromial artery. This sturdy artery originates from the axillary artery, a major vessel supplying the upper limb. The pectoral branch courses along the undersurface of the

pectoralis major muscle, sending perforating branches that penetrate the muscle and nourish the overlying skin and subcutaneous tissues. This creates a rich vascular network that supports the flap's viability even when it is transposed to a distant recipient site. In addition to the primary supply, the pectoralis major flap benefits from secondary blood supply from the internal mammary artery and lateral thoracic artery. These collateral vessels provide an additional layer of vascular support, ensuring flap survival even in situations where the primary blood supply may be compromised due to factors such as anatomical variations, previous surgery, or radiation therapy. This robust and multi-layered blood supply is crucial for flap survival, especially in the context of head and neck reconstruction. The head and neck region is characterized by complex anatomy and a high density of vital structures. Reconstructive procedures in this

area often involve extensive dissection and manipulation of tissues, which can potentially compromise blood flow to the flap. The pectoralis major flap's inherent vascularity makes it a reliable option, even in challenging reconstructive scenarios. The anatomical location of the pectoralis major muscle on the anterior chest wall contributes significantly to the ease of flap harvest. This accessibility simplifies the surgical procedure, reduces operative time, and minimizes potential complications at the donor site. The pectoralis major muscle is situated superficially on the chest, making it readily accessible through a relatively straightforward incision. This allows for efficient flap elevation with minimal disruption to surrounding tissues and vital structures. The surgical technique for harvesting the pectoralis major flap is well-established and relatively straightforward. This makes it a technically less demanding option compared to some other flaps used in head and neck reconstruction, which may require more complex dissection or microvascular anastomosis. The flap can be harvested with or without a portion of the underlying rib, depending on the specific reconstructive needs. The ease of harvest translates to several advantages, including reduced operative time, minimized blood loss, and potentially fewer complications at the donor site. This contributes to an improved overall patient experience and facilitates a smoother postoperative recovery. The pectoralis major flap's versatility extends beyond its vascular reliability and ease of harvest. It offers the capacity to provide substantial tissue bulk, making it particularly well-suited for reconstructing extensive or complex defects in the head and neck. The muscle itself provides a significant volume of well-vascularized tissue, which can be used to fill large cavities, restore lost contour, and provide structural support. The overlying skin and subcutaneous tissues add further bulk and can be tailored to match the specific needs of the defect. This ample tissue volume allows for the restoration of natural contours and facilitates the functional rehabilitation of critical structures in the head and neck region. For example, in the case of oropharyngeal

reconstruction, the pectoralis major flap can be used to reconstruct the soft palate, posterior pharyngeal wall, and base of tongue, restoring swallowing function and speech intelligibility. The ability to provide substantial tissue bulk makes the pectoralis major flap particularly valuable in complex reconstructive scenarios. For instance, in cases of extensive tumor resection or traumatic injuries, where significant tissue loss has occurred, the flap can effectively fill the defect and restore the structural integrity of the head and neck. The pectoralis major flap has proven its efficacy in addressing a wide spectrum of head and neck defects, showcasing its versatility in managing various reconstructive challenges. In the context of oncological reconstruction, the pectoralis major flap is frequently employed following the resection of tumors in the oral cavity, oropharynx, hypopharynx, and larynx. Its ability to provide ample tissue bulk and restore critical anatomical structures makes it a valuable tool in restoring both form and function after tumor ablation. The pectoralis major flap has also found applications in the reconstruction of traumatic defects, such as those resulting from gunshot wounds, motor vehicle accidents, and burns. Its ability to provide well-vascularized tissue coverage and restore lost contour makes it an effective option for managing these challenging injuries. Furthermore, the pectoralis major flap can be used to address congenital defects, such as cleft palate and micrognathia. Its versatility in providing both soft tissue and bony support makes it a valuable tool in correcting these complex anomalies. While the pectoralis major flap offers numerous advantages, it is essential to consider potential donor site morbidity. Careful patient selection, meticulous surgical technique, and vigilant postoperative monitoring can mitigate these risks. The most common complications include seroma formation (fluid accumulation), hematoma (blood collection), wound dehiscence (wound breakdown), and sensory changes in the chest and axilla. These complications are generally minor and can be managed conservatively. Careful patient selection, meticulous

surgical technique, and vigilant postoperative monitoring can minimize the risk of donor site complications. In most cases, donor site morbidity is minimal and well-tolerated by patients. The advent of free tissue transfer techniques has expanded the reconstructive options available for head and neck defects. However, the pectoralis major flap remains a valuable tool in the reconstructive surgeon's armamentarium. Free flaps, which involve the transfer of tissue from a distant donor site with microvascular anastomosis, offer greater versatility in terms of tissue selection and can be used to reconstruct even the most complex defects. Despite the advancements in free tissue transfer, the pectoralis major flap remains a relevant and frequently used option for head and neck reconstruction. Its robust vascularity, ease of harvest, and ample tissue bulk make it a suitable choice in many scenarios, particularly when simpler reconstructive options are preferred or when free tissue transfer is not feasible. The choice of flap should always be individualized based on the specific characteristics of the defect, the patient's overall health, and the surgeon's experience and preference. In many cases, the pectoralis major flap continues to be a reliable and effective option for restoring form and function in the head and neck region.¹¹⁻¹³

In the case described, the patient's history of parotid gland cancer and prior treatment with chemotherapy, radiotherapy, and embolization posed additional challenges to the reconstructive effort. These prior interventions can induce tissue fibrosis, compromise vascularity, and increase the risk of wound healing complications. The inherent resilience of the pectoralis major flap, with its robust vascular supply, makes it a suitable option for reconstructing defects in patients with prior treatment history. The flap's ability to withstand the challenges posed by irradiated or scarred tissue contributes to its success in these complex scenarios. Cancer treatment modalities, such as chemotherapy, radiotherapy, and embolization, while aimed at eradicating the malignancy, can have detrimental effects on the surrounding healthy tissues. These effects can pose

significant challenges in the context of reconstructive surgery. Chemotherapy drugs are systemic agents designed to kill rapidly dividing cancer cells. However, they can also affect healthy tissues, particularly those with high turnover rates, such as bone marrow, hair follicles, and the lining of the digestive tract. In the context of head and neck reconstruction, chemotherapy can impair wound healing by disrupting cellular processes involved in tissue repair and regeneration. It can also lead to decreased vascularity, making the tissues more susceptible to ischemia and necrosis. Radiation therapy utilizes high-energy rays to target and destroy cancer cells. However, the surrounding healthy tissues can also be affected by the radiation, leading to a range of complications. One of the most significant concerns is radiation-induced fibrosis, a process where the irradiated tissues become scarred and less pliable. This fibrosis can impair blood flow, reduce tissue oxygenation, and hinder wound healing. Embolization is a procedure that involves the deliberate blockage of blood vessels to cut off the blood supply to a tumor. While effective in shrinking the tumor, it can also compromise the vascularity of surrounding tissues. This can make it challenging to perform reconstructive surgery, as the compromised blood flow may impair the survival of the flap. The challenges posed by prior cancer treatment can significantly complicate reconstructive efforts. The compromised tissue health can increase the risk of flap failure, infection, and wound breakdown. The decreased vascularity in previously treated tissues can impair the survival of the flap, as it may not receive adequate blood supply to support its metabolic needs. This can lead to flap necrosis, a serious complication that can necessitate further surgery and delay the healing process. Radiation-induced fibrosis can make it difficult to create a healthy recipient site for the flap. The scarred and fibrotic tissue may lack the pliability and vascularity necessary for successful flap integration. This can increase the risk of flap failure and wound complications. The impaired wound healing capacity of previously treated tissues can increase the risk of

wound breakdown and infection. This can delay the healing process, prolong hospital stays, and compromise the overall outcome of the reconstruction. Despite these challenges, the pectoralis major flap has demonstrated its resilience and effectiveness in reconstructing defects in patients with prior cancer treatment history. Its inherent characteristics make it well-suited to withstand the challenges posed by compromised tissue health. The robust vascularity of the pectoralis major flap, with its primary and secondary blood supply, provides a significant advantage in reconstructing previously treated tissues. The flap's ability to maintain adequate blood flow, even in the setting of decreased vascularity, enhances its chances of survival and successful integration. The pectoralis major flap has also shown a degree of tolerance to the effects of radiation therapy. While radiation can cause tissue damage, the flap's inherent resilience allows it to withstand some degree of radiation-induced changes and still maintain its viability. This tolerance is attributed to the flap's rich blood supply and the ability of the muscle to regenerate even after radiation exposure. The flap's ability to conform to irregular recipient sites and integrate with scarred tissue further contributes to its success in reconstructing previously treated defects. The pliable nature of the flap allows it to adapt to the contours of the recipient site, even in the presence of fibrosis and scar tissue. While the pectoralis major flap offers a resilient option for reconstructing previously treated defects, meticulous surgical planning and execution are crucial for optimizing outcomes. Careful patient selection is essential to ensure that the pectoralis major flap is the most appropriate reconstructive option. Factors such as the extent of prior treatment, the degree of tissue damage, and the patient's overall health should be carefully considered. In some cases, where the tissue damage is extensive or the patient's health is compromised, alternative reconstructive options may be more suitable. Meticulous surgical technique is paramount for minimizing complications and achieving the best possible outcomes. This includes careful handling of

tissues, strict adherence to aseptic principles, and precise flap design and inset. The surgeon's expertise and experience play a crucial role in ensuring the success of the reconstruction. Adequate preparation of the recipient site is crucial for successful flap integration. This may involve debridement of unhealthy tissue, release of scar contractures, and meticulous hemostasis to ensure a healthy and well-vascularized recipient bed. Proper recipient site preparation enhances the chances of flap survival and reduces the risk of complications. Vigilant postoperative monitoring is essential for early detection and management of potential complications. This includes close observation of the flap for signs of vascular compromise, such as color changes, coolness, or decreased capillary refill. It also involves monitoring for signs of infection, such as redness, swelling, pain, or fever. Early detection and prompt intervention can significantly improve outcomes. In some cases, adjuvant therapies, such as hyperbaric oxygen therapy (HBOT), may be considered to enhance wound healing and improve the success of the reconstruction. HBOT involves breathing 100% oxygen in a pressurized chamber, which increases the amount of oxygen dissolved in the blood. This increased oxygenation can promote angiogenesis (formation of new blood vessels), enhance tissue regeneration, and reduce the risk of infection.^{14,15}

The design and execution of the pectoralis major flap require careful consideration of the specific characteristics of the defect being addressed. Factors such as the size, location, and depth of the defect, as well as the patient's individual anatomy, influence the surgical planning process. In this case, the large size of the defect, measuring 8 x 12 cm, necessitated a flap with sufficient dimensions to ensure adequate coverage and restore the natural contours of the neck. The flap was meticulously designed and harvested to match the defect's dimensions precisely, ensuring optimal aesthetic and functional outcomes. The success of any reconstructive procedure hinges on meticulous preoperative planning. This is particularly crucial when utilizing the pectoralis major flap, as the

flap design and execution must be tailored to the specific characteristics of the defect and the patient's individual anatomy. The first step in preoperative planning is a thorough assessment of the defect. This involves a detailed evaluation of its size, location, and depth, taking into account the involvement of surrounding structures. The reconstructive surgeon must also consider the potential impact of the defect on critical functions such as speech, swallowing, and breathing. For instance, a defect involving the oral cavity may affect speech articulation and swallowing, while a defect in the larynx may compromise airway patency. In addition to evaluating the defect, the surgeon must also assess the patient's overall health, nutritional status, and any comorbidities that may influence the reconstructive process. Factors such as previous cancer treatment, smoking history, and diabetes can affect wound healing and flap viability. Patients who have undergone radiation therapy may have compromised tissue vascularity and healing capacity, while those with diabetes may be more susceptible to infections. Imaging studies, such as computed tomography (CT) scans and magnetic resonance imaging (MRI), play a crucial role in preoperative planning. These studies provide detailed information about the defect's extent, its relationship to surrounding structures, and the availability of recipient vessels for flap anastomosis. For example, a CT scan can help visualize the bony structures and assess the involvement of the mandible or maxilla, while an MRI can provide detailed information about the soft tissues and identify any potential areas of compromised vascularity. In complex cases, virtual surgical planning (VSP) may be employed to create a three-dimensional model of the defect and surrounding anatomy. This allows the surgeon to simulate the reconstructive procedure and optimize flap design and placement before entering the operating room. VSP can help identify potential challenges and improve the accuracy of flap design, leading to better aesthetic and functional outcomes. The design of the pectoralis major flap must be tailored to the specific needs of the defect. Factors

such as the defect's size, location, and depth, as well as the patient's individual anatomy, influence the flap design process. The flap must be large enough to provide adequate coverage of the defect and restore the natural contours of the head and neck. In the case described, the large size of the defect (8 x 12 cm) necessitated a flap with sufficient dimensions to ensure complete closure and avoid wound tension. If the flap is too small, it may not provide adequate coverage, leading to wound dehiscence or inadequate functional restoration. The skin paddle, which is the portion of the flap that will be visible after the reconstruction, must be carefully designed to match the color, texture, and thickness of the surrounding skin. The surgeon must also consider the location of the skin paddle on the chest to minimize donor site morbidity and optimize the aesthetic outcome. For example, if the defect is located in the lower face, the skin paddle may be harvested from the lower chest to achieve a better color and texture match. The amount of muscle included in the flap depends on the reconstructive needs. In some cases, only a portion of the pectoralis major muscle may be required, while in others, the entire muscle may be harvested. The vascular pedicle, which supplies blood to the flap, must be carefully preserved during flap elevation to ensure its viability. The length and diameter of the pedicle should be assessed to ensure adequate blood flow to the flap. The orientation of the flap is crucial for achieving optimal aesthetic and functional outcomes. The flap must be positioned to restore the natural contours of the head and neck and avoid creating unsightly bulges or depressions. The inset of the flap, which involves suturing it into place, must be performed meticulously to ensure proper alignment and tension. The surgeon must also consider the potential impact of flap placement on surrounding structures, such as nerves and blood vessels. While preoperative planning provides a roadmap for the reconstructive procedure, the surgeon must be prepared to adapt to the intraoperative findings. The surgical field may reveal unexpected challenges that require modifications to the flap design or execution.

The actual defect encountered during surgery may differ slightly from the preoperative assessment. The surgeon must be prepared to adjust the flap design or harvest additional tissue if necessary to ensure adequate coverage. For example, if the defect is larger than anticipated, the surgeon may need to extend the flap or harvest a second flap to achieve complete closure. The vascularity of the recipient site and the flap's pedicle must be carefully assessed intraoperatively. If the blood supply is compromised, the surgeon may need to consider alternative reconstructive options or perform microvascular anastomosis to ensure flap survival. Doppler ultrasound can be used to assess blood flow in the flap's pedicle and recipient vessels. Meticulous hemostasis (control of bleeding) is crucial to prevent hematoma formation and promote wound healing. The wound closure must be performed in layers, using appropriate suture techniques to minimize tension and optimize the aesthetic outcome. The surgeon must also consider the placement of drains to prevent fluid accumulation and promote wound healing. Postoperative care plays a crucial role in ensuring the success of the reconstruction. The patient must be closely monitored for signs of flap complications, such as vascular compromise, infection, or wound breakdown. The flap should be regularly assessed for color, temperature, capillary refill, and turgor. Any signs of vascular compromise should be addressed promptly to prevent flap necrosis. The patient's vital signs, including heart rate, blood pressure, and oxygen saturation, should also be closely monitored. Proper wound care is essential to prevent infection and promote healing. This may involve regular dressing changes, wound irrigation, and the use of topical or systemic antibiotics. The patient should be instructed on proper wound care techniques and advised to avoid activities that may disrupt the healing process. The patient should be educated about the signs and symptoms of flap complications and instructed to report any concerns to the healthcare team immediately. This includes information about changes in flap color, temperature, or sensation, as well as

signs of infection, such as redness, swelling, or discharge.¹⁶⁻¹⁸

The success of head and neck reconstruction hinges not only on the technical aspects of flap design and execution but also on a comprehensive approach to patient care. This includes thorough preoperative evaluation, meticulous surgical technique, and vigilant postoperative monitoring. Preoperative assessment of the patient's overall health, nutritional status, and any comorbidities is crucial for minimizing surgical risk and optimizing postoperative recovery. Clear communication with the patient regarding treatment options, potential risks, and expected outcomes ensures informed decision-making and fosters a collaborative approach to care. Intraoperatively, meticulous surgical technique is paramount for minimizing complications and achieving the best possible outcomes. Careful handling of tissues, strict adherence to aseptic principles, and precise flap design and inset are essential for ensuring flap viability and promoting wound healing. Postoperative care plays a pivotal role in ensuring the success of the reconstruction. Close monitoring of the patient's vital signs, flap viability, and wound healing allows for early detection and management of potential complications. Adequate pain control, nutritional support, and appropriate wound care contribute to the patient's comfort and facilitate an uneventful recovery. The preoperative period provides a crucial window of opportunity to optimize the patient's health, assess their individual needs, and prepare them for the challenges of head and neck reconstruction. A comprehensive preoperative evaluation, combined with clear communication and patient education, sets the stage for a successful outcome. A detailed medical history, including any prior cancer treatment, surgeries, or medical conditions, is essential to identify potential risk factors and tailor the treatment plan accordingly. A thorough physical examination, including assessment of the patient's nutritional status and any signs of infection or inflammation, is also crucial. Adequate nutrition is essential for wound healing and

overall recovery. Patients with malnutrition or those at risk of malnutrition should receive nutritional support in the preoperative period to optimize their metabolic reserves and enhance their ability to withstand the stress of surgery. Patients with comorbidities, such as diabetes, hypertension, or cardiovascular disease, require careful management in the preoperative period to minimize the risk of complications. This may involve optimizing their medication regimen, controlling their blood sugar levels, and ensuring adequate hydration. Head and neck reconstruction can be a physically and emotionally challenging experience for patients. Providing psychological support and counseling can help patients cope with the stress of surgery and prepare for the recovery process. Clear communication with the patient regarding treatment options, potential risks, and expected outcomes is crucial for ensuring informed consent and fostering a collaborative approach to care. The patient should be actively involved in the decision-making process and feel empowered to express their concerns and preferences. The intraoperative period demands meticulous surgical technique and attention to detail to minimize complications and achieve the best possible reconstructive outcomes. Careful handling of tissues, strict adherence to aseptic principles, and precise flap design and inset are essential for ensuring flap viability and promoting wound healing. Gentle tissue handling is crucial to minimize trauma and preserve tissue vascularity. The surgeon must avoid excessive traction, compression, or crushing of tissues, as these can compromise blood flow and impair wound healing. Maintaining a sterile surgical field is paramount to prevent infection, a serious complication that can compromise flap viability and delay healing. The surgical team must adhere to strict aseptic techniques, including proper hand hygiene, gowning, and gloving, as well as the use of sterile instruments and drapes. The flap must be meticulously designed and harvested to match the defect's dimensions and ensure adequate coverage. The inset of the flap, which involves suturing it into place, must be performed with precision to avoid

tension, distortion, or compression of the vascular pedicle. Achieving meticulous hemostasis (control of bleeding) is crucial to prevent hematoma formation, which can compromise flap viability and increase the risk of infection. The surgeon must also carefully manage any dead space (empty pockets) in the wound to prevent fluid accumulation and promote healing. The postoperative period requires vigilant monitoring and proactive management of potential complications to ensure the success of the reconstruction. Close observation of the patient's vital signs, flap viability, and wound healing allows for early detection and intervention, minimizing morbidity and optimizing outcomes. Close monitoring of the patient's vital signs, including heart rate, blood pressure, respiratory rate, and oxygen saturation, is crucial to detect any signs of instability or complications. Any significant deviations from baseline should be promptly investigated and managed. The flap should be regularly assessed for signs of vascular compromise, such as color changes, coolness, decreased capillary refill, or decreased turgor. Early detection of vascular compromise allows for prompt intervention, such as flap exploration or revision, to salvage the flap and prevent necrosis. Proper wound care is essential to prevent infection and promote healing. This may involve regular dressing changes, wound irrigation, and the use of topical or systemic antibiotics. The patient should be instructed on proper wound care techniques and advised to avoid activities that may disrupt the healing process. Adequate pain control is crucial for the patient's comfort and facilitates early mobilization and rehabilitation. The surgeon should prescribe appropriate pain medications and monitor the patient's pain levels regularly. Maintaining adequate nutrition is essential for wound healing and overall recovery. Patients may require nutritional support in the postoperative period, either orally or through intravenous fluids, to ensure adequate caloric and protein intake. The patient should be educated about the signs and symptoms of flap complications and instructed to report any concerns to the healthcare team immediately. Regular follow-up

appointments are essential to monitor the patient's progress, assess flap viability, and address any complications that may arise.^{19,20}

4. Conclusion

The pectoralis major flap has proven to be a reliable and versatile option for reconstructing head and neck defects, offering several advantages including robust vascularity, ease of harvest, and the ability to provide substantial tissue bulk. Its versatility has been demonstrated across a range of clinical scenarios, including tumor resection, trauma reconstruction, and congenital defect correction. The flap's resilience and ability to withstand the challenges posed by compromised tissue health make it a valuable option for patients with prior cancer treatment history. Careful preoperative planning, meticulous surgical technique, and vigilant postoperative monitoring are essential for optimizing outcomes and minimizing complications. Thorough patient assessment, clear communication, and informed consent contribute to a patient-centered approach to care. Intraoperative attention to detail, including gentle tissue handling, strict adherence to aseptic principles, and precise flap design and inset, ensures flap viability and promotes wound healing. Postoperative care focuses on close monitoring of the patient's vital signs, flap viability, and wound healing, allowing for early detection and management of potential complications.

5. References

1. Wang C-H, Wong Y-K, Wang C-P, Wang C-C, Jiang R-S, Lai C-S, et al. Risk factors of recipient site infection in head and neck cancer patients undergoing pectoralis major myocutaneous flap reconstruction. *Eur Arch Otorhinolaryngol*. 2015; 272(11): 3475–82.
2. Zhang W, Li M, Liu N, Chen F, Meng Q. The application of double pedicle blood supply pectoralis major myocutaneous flap in reconstruction of laryngeal and hypopharyngeal defect after carcinoma resection. *Lin Chuang Er Bi Yan Hou Tou Jing Wai Ke Za Zhi*. 2015; 29(7): 589–92.
3. Chen J, Huang W, Li Z, Zhou X, Yu J, Bao R, et al. Modified pectoralis major myocutaneous flap in reconstruction of head and neck defects. *Zhonghua Er Bi Yan Hou Tou Jing Wai Ke Za Zhi*. 2015; 50(5): 388–91.
4. Zhang W, Li M, Liu N, Chen F, Meng Q. The application of double pedicle blood supply pectoralis major myocutaneous flap in reconstruction of laryngeal and hypopharyngeal defect after carcinoma resection. *Lin Chuang Er Bi Yan Hou Tou Jing Wai Ke Za Zhi*. 2015; 29(7): 589–92.
5. Chen J, Huang W, Li Z, Zhou X, Yu J, Bao R, et al. Modified pectoralis major myocutaneous flap in reconstruction of head and neck defects. *Zhonghua Er Bi Yan Hou Tou Jing Wai Ke Za Zhi*. 2015; 50(5): 388–91.
6. Chen W-L, Wang Y-Y, Zhang D-M, Fan S, Lin Z-Y. Extended vertical lower trapezius island myocutaneous flap versus pectoralis major myocutaneous flap for reconstruction in recurrent oral and oropharyngeal cancer. *Head Neck*. 2016; 38(Suppl 1): E159-64.
7. Refos JWJ, Witte BI, de Goede CJT, de Bree R. Shoulder morbidity after pectoralis major flap reconstruction. *Head Neck*. 2016; 38(8): 1221–8.
8. Knopf A, Mansour N, Hofauer B, Bier H, Scherer EQ. Outcome after pharyngeal reconstruction using pectoralis major and radial forearm flap after resection of pharyngeal and laryngeal squamous cell carcinomas. *Eur Arch Otorhinolaryngol*. 2016; 273(9): 2637–42.
9. Nguyen S, Thuot F. Functional outcomes of fasciocutaneous free flap and pectoralis major flap for salvage total laryngectomy. *Head Neck*. 2017; 39(9): 1797–805.
10. Pradhan P, Samal S, Preetam C, Samal DK, Parida PK. Pectoralis major myocutaneous flap reconstruction for the mandibular defects

- in advanced oral cavity malignancies: a retrospective study of 30 cases. *Indian J Otolaryngol Head Neck Surg.* 2018; 70(3): 415–20.
11. Pradhan P, Samal S, Preetam C. Pectoralis major myocutaneous flap for the reconstruction of the palatal defect. *Indian J Otolaryngol Head Neck Surg.* 2019; 71(1): 132–5.
 12. Chu Y-H, Lai W-S, Lin Y-Y, Liu S-C, Lee J-C. Pharyngeal reconstruction using a U-shaped pectoralis major myocutaneous flap: an effective technique that should not be forgotten. *Eur Arch Otorhinolaryngol.* 2020; 277(1): 217–20.
 13. Deo RP, Kumar BA, Sagayaraj, Mohiyuddin A. Utility of island bipaddle pectoralis major myocutaneous flap in reconstruction of full thickness lower gingivo-buccal sulcus defects. *Int J Otorhinolaryngol Head Neck Surg.* 2020; 7(1): 56.
 14. Sahu PK, Kumar S. Bipaddle pectoralis major myocutaneous flap for single stage reconstruction of oromandibular defects. *Indian J Otolaryngol Head Neck Surg.* 2020; 72(1): 44–8.
 15. Shanmugam S, Hussain SA, Michael R. Factors increasing the risk of flap complications following pectoralis major myocutaneous flap reconstruction for head and neck cancers. *Int J Otorhinolaryngol Head Neck Surg.* 2020; 6(6): 1138.
 16. Singh A, Subash A, Majumdar K, Bhinyaram. Cysticercosis infested pectoralis major myocutaneous flap reconstruction: An ethical dilemma. *Indian J Otolaryngol Head Neck Surg.* 2022; 74(Suppl 2): 2231–5.
 17. Dhiwakar M, Shanmugam J, Khan ZA, Mehta S, Karthik K. Distal positioning of the skin paddle of pectoralis major myocutaneous flap in head and neck reconstruction. *Head Neck.* 2023; 45(11): 2819–28.
 18. Mitra S, Panda S, Amit Singh C, Mani S, Thakar A. Folded pectoralis major myocutaneous flap for reconstruction of complex oral commissural defects: a retrospective case series and technical report. *Indian J Otolaryngol Head Neck Surg.* 2024; 76(1): 1251–4.
 19. Judd RT, McCrary HC, Farlow JL, Li M, Godsell J, Kneuert PJ, et al. Pedicled osteomyocutaneous pectoralis major flap with osseous rib harvest for salvage mandibular reconstruction: Case and technique. *Head Neck.* 2024; 46(2): 447–51.
 20. Mahajan A, Kumar JA, Singh AK, Sharma NK, Durrani F, Agrawal NK, et al. Quality of life assessment and comparative analysis of anterolateral thigh flap and pectoralis major myo-cutaneous flap for reconstruction of post excision defects of oral squamous cell carcinoma. *Indian J Otolaryngol Head Neck.* 2024; 76(4): 3390–5.